

# **EXHIBIT B**

**BEST AVAILABLE COPY**

Journal of Drug Monitoring  
2004;17: 373-377 © 1998 Lippincott Williams & Wilkins, Philadelphia

## Pharmacodynamics, Pharmacokinetics, and Therapeutic Drug Monitoring of Glycopeptides

Alasdair P. MacGowan

Bristol Centre for Antimicrobial Research and Evaluation, Southmead Health Services, NHS Trust; and University of Bristol, Department of Medical Microbiology, Southmead Hospital, Westbury-on-Trym, United Kingdom

**Summary:** The glycopeptide antibacterial drugs, vancomycin and teicoplanin, are widely used in hospitals for therapy of severe or multiresistant infection that has a positive result on Gram's stain test. Although vancomycin resistance is common in some hospital-acquired *Enterococcus* sp and resistance to teicoplanin occurs among *Staphylococcus* sp glycopeptides remain the cornerstone of therapy for infection due to methicillin-resistant *Staphylococcus aureus* (MRSA), coagulase-negative *Staphylococcus* organisms, and infection related to implanted devices. Therapeutic drug monitoring (TDM) of these agents remains controversial, but advances in our understanding of their pharmacodynamics and further clinical studies are helping clarify the situation. In the future, a more rational approach to monitoring will probably result in less intensive monitoring of vancomycin but more intensive monitoring of teicoplanin. **Key Words:** Vancomycin—Teicoplanin—Pharmacodynamics—Therapeutic drug monitoring.

Pharmacodynamics offers an opportunity to relate knowledge about an antimicrobial drug's *in-vitro* susceptibility, minimum inhibitory concentration (MIC), post-antibiotic effect (PAE), pattern of bactericidal action, and interactions with immune cells to pharmacokinetics to optimize drug dosing regimens.

Postantibiotic effect is the ability of an antibacterial to suppress the generation of bacteria for several hours after antibacterial concentrations have fallen below the MIC. Although the exact mechanism of the PAE is unknown, it may be related to repair of damaged, but not killed, cells; separation of bound drug from target; or synthesis of new enzymes or proteins (1). Penicillins, cephalosporins, macrolides, and aminoglycosides have a PAE against bacteria that have a positive result on Gram's stain tests (2). Although the measurement of the PAE depends on the method used (3), vancomycin has been

shown by a number of techniques to have a PAE of 2 to 3 hours against *Staphylococcus aureus* (2,4,5).

Teicoplanin also has a PAE that appears to be longer than that of vancomycin (6,7). If a concentration well below the MIC is allowed to remain instead of completely removing the antibiotic during the measurement of the PAE, the PAE duration is doubled for vancomycin, which is termed a sub-MIC effect. This measure is akin to the exponential decay of a drug concentration in serum (8).

Increasing the concentration of vancomycin in the therapeutic range (i.e., from 5 mg/L to 40 mg/L) does not increase the time to kill 99.9% of the bacterial population or the rate of kill (9); the rate of killing is slower for teicoplanin than for vancomycin, perhaps because of the former's high protein binding (10).

The pharmacodynamics of glycopeptides studied in several animal models support the concept that high initial concentrations offer no advantage in bacterial killing or mortality, whereas higher, sustained concentrations or more frequent dosing have improved survival in animal models of infective endocarditis (11,12).

In a complex analysis using a mouse model, multiple

Received February 11, 1998; accepted July 23, 1998.  
Address correspondence and reprint requests to Alasdair P. MacGowan, University of Bristol, Department of Medical Microbiology, Southmead Hospital, Westbury-on-Trym, Bristol BS10 5NB, United Kingdom.

pharmacodynamic parameters were compared with the effective dose  $_{50}$  ( $ED_{50}$ ). The time serum concentration (T) exceeded the MIC ( $T > MIC$ ) and was best related to  $ED_{50}$  when treating penicillin-resistant *Pneumococci* organisms with either vancomycin or teicoplanin (13). Use of an in vitro, continuous bacterial culture model supports this finding. By simulating four different therapeutic regimens with various peak or trough concentrations or areas under the curve (AUC), but maintaining  $T > MIC$  at 100%, it was shown that there were no differences in the degree or rate of *S. aureus* killing (14).

Laboratory and animal evaluation of glycopeptide pharmacodynamics indicates that glycopeptides do not show concentration-dependent killing in the therapeutic range; hence, high peak concentrations are unlikely to be of benefit. In addition, they have a PAE and sub-MIC effects, indicating that serum concentrations need not exceed the MIC for all of the dosing interval and that  $T > MIC$  or sustained concentrations are related to outcome. Protein binding affects bacterial killing with teicoplanin. Therefore, the dosing interval is probably best optimized as  $T > MIC$  plus PAE, although clinically, with conventional doses of vancomycin,  $T > MIC$  is 100%. This result had led some to propose that smaller doses than the standard 2 grams per day of vancomycin may be just as effective in clinical practice; alternatively, longer dosing intervals may be appropriate for glycopeptides (15).

### PHARMACOKINETICS

The pharmacokinetics of vancomycin and teicoplanin have been extensively studied and are known to vary in different patient groups. For example, vancomycin handling is changed in renal impairment (16-18); obesity (19,28), liver failure (21), various renal support therapies (22-27), neutropenia (28), malignancy (29), age, and gender (30); and with sepsis and its therapy (Table 1 [31,32]).

Similarly, teicoplanin pharmacokinetics are altered in renal impairment (33,34), renal support therapies (35,36); children and the elderly (37), intravenous (IV) drug abusers (38), burn patients (9), and neutropenia (see Table 1 [40]). In addition, it is clear that standard dosing of teicoplanin (400 mg  $\times$  2 for 24 hours, then 400 mg 24 hourly) results in significant numbers of patients having predose serum concentrations of 10 mg/L (40).

However, pharmacokinetic variability on its own can rarely be a justification for TDM and only becomes important if serum concentrations can be linked to toxicity or efficacy. This link has been tenuous for glycopeptides and continued TDM was questioned in the late 1980s and

TABLE 1. Patient factors affecting vancomycin or teicoplanin pharmacokinetics

Patient Factor	Pharmacokinetic change	Reference no.
<b>Vancomycin</b>		
Renal impairment	Increasing t <sub>1/2</sub> with decreasing creatinine clearance	17
Chronic intermittent hemodialysis	As for renal impairment: little drug removed by dialysis	22
Chronic intermittent peritoneal dialysis	Prolonged t <sub>1/2</sub>	23
Continuous veno-venous hemofiltration or ultrafiltration	Increased clearance compared to hemodialysis	25,27
Obesity	Shorter t <sub>1/2</sub> , larger volume of distribution	18,19
Liver failure	Longer t <sub>1/2</sub>	21
Age	Longer t <sub>1/2</sub> in neonates than infants and children	29,36
Sepsis	Prolonged t <sub>1/2</sub>	23
<b>Teicoplanin</b>		
Renal impairment	Increasing t <sub>1/2</sub> with decline in creatinine clearance	33,34
Chronic intermittent hemodialysis	Reduced clearance	35
Chronic ambulatory peritoneal dialysis	Prolonged t <sub>1/2</sub> , increased volume of distribution	32
Continuous veno-venous hemofiltration	Prolonged t <sub>1/2</sub>	33
Continuous hemofiltration	t <sub>1/2</sub> not prolonged	36
Intravenous drug abusers	Increased clearance	38
Burn patients	Increased t <sub>1/2</sub>	39
Neutropenia	Increased elimination, larger interindividual variability	40

early 1990s (42-46). This process has now resulted in the emergence of new data.

### TOXICITY

Vancomycin serum monitoring, if performed, is aimed at reducing the risks of nephrotoxicity or ototoxicity: it will not reduce immediate or infusion-related toxicities. Ototoxicity is difficult to assess clinically and data is sketchy because they are often composed of case reports in patients with renal failure, who sometimes have high serum concentrations. It is not sufficient to make any association between concentrations and toxicity.

The incidence of nephrotoxicity is probably less than 5% in patients treated with vancomycin alone, but higher if a combination of vancomycin plus an aminoglycoside is used (47). Toxicity is also associated with longer courses of therapy and the original report of Faber and Moellering (47) linked three patients to trough concentrations of 30 mg/L to 65 mg/L before the onset of tox-

## PHARMACODYNAMICS, PHARMACOKINETICS, AND TDM OF GLYCOPOLYPTIDES

475

icity. There are now several reports that vancomycin TDM services involving prescription review, blood concentration measurement, and dose modification by clinical pharmacists can reduce the incidence of nephrotoxicity. In a 1994 prospective cohort study in a teaching hospital, 116 patients who received more than 4 days' therapy and were not neutropenic or in an intensive care unit or established renal failure were studied. It was shown that the rate of nephrotoxicity was 24% in patients not randomized to the TDM service, compared with 7% of those who received TDM (43). A similar and prospective randomized study in 70 patients with hematologic malignancy indicated that nephrotoxicity was lower in patients recruited into the TDM arm (mild toxicity, 13.5%; moderate, 0%) compared with those who received no TDM (mild, 33%; moderate, 9.1% [49]). Furthermore, in a retrospective review of 273 patients with positive results of infection with Gram's stain, it was shown that serum vancomycin concentrations before onset of nephrotoxicity were higher (23.5 [2.5 mg/L]) in those in whom toxicity developed than in those in whom it did not (10.2 [3.8 mg/L] [50]). In contrast, in a prospective study of patients randomized to have dose adjustment to achieve predose concentrations in the ranges of 5 mg/L to 10 mg/L, 10 mg/L to 15 mg/L, or 15 mg/L to 25 mg/L, no correlation was found to nephrotoxicity (51).

Thrombocytopenia associated with large doses of teicoplanin (30 mg/kg per day) has recently been related to trough concentrations; for those with trough concentrations of more than 60 mg/L, eight of 58 patients had a decrease in platelets, whereas with trough concentrations of less than 60 mg/L, 12 of 251 had a decrease in platelets ( $p < 0.05$ ) [52]).

## OUTCOME

Retrospective data reviews of teicoplanin clinical trials have indicated that serum concentrations are related to clinical outcomes. In an open multicenter study in which most patients had right-sided infection due to *S. aureus* and predose concentrations had been adjusted to between 10 mg/L to 15 mg/L, it was reported that postdose concentrations of teicoplanin of more than 40 mg/L were associated with improved outcome (53). A further study (mainly of bone infection due to *S. aureus*) suggested that larger doses than were conventionally used at that time were required for successful therapy and that average troughs were 36.3 ( $n = 10$ ) in those successfully treated, and only 9.7 mg/L in the three clinical failures (54). A retrospective review of three trials in the United States indicated an association between increased dose, high trough serum levels, trough concentration/MIC ratio

and days to clear bacteremia, fever days, and clinical improvement (55). A further retrospective review of 58 cases published with sufficient pharmacokinetic and susceptibility data for analysis indicated a relationship between predose and postdose teicoplanin concentrations and predose/MIC or postdose/MIC ratios and clinical outcomes. Dose was not related to outcome in this review, in which most of the patients had severe staphylococcal infection (56). In a more recent review of 92 patients with *S. aureus* bacteremia using a multivariate analysis to relate age, weight, dose, loading dose, combination therapy, and serum concentrations to outcome has shown that only trough concentration and age were significantly related to outcome (57). A prospective study of teicoplanin to treat *S. aureus* infective endocarditis showed that if the predose concentration was less than 20 mg/L, six of 10 patients failed, compared with one of 11 if the concentration was more than 20 mg/L ( $p < 0.05$ ) (58).

The data relating to vancomycin serum concentration to efficacy is less clear. Two prospective studies showed that therapeutic drug monitoring (TDM) services had no effect on efficacy (48,49) and an intervention in which patients were deliberately stratified into three groups with predose targets of 5 mg/L to 10 mg/L, 10 mg/L to 15 mg/L, or 15 mg/L to 25 mg/L showed no difference in fever days or clinical outcome (51). In contrast, two retrospective reviews were able to relate serum concentrations to outcome measures. In a retrospective review of 273 patients with positive results of infection proven with Gram's stain, Zimmerman and colleagues (50) were able to relate troughs of more than 10 mg/L to a reduced number of fever days and an improved white blood cell response but not to lengths of stay or mortality. Mulhern and colleagues (59) related trough concentrations to relapse rates in patients with peritonitis who were treated with continuous ambulatory peritoneal dialysis for end-stage renal disease. When the mean predose concentration was less than 12 mg/L, 9 of 14 patients relapsed; when it was more than 12 mg/L, none of 17 relapsed.

In conclusion, pharmacodynamic principles indicate that predose glycopeptides should be related to the outcome of infection measures. Evidence now exists in humans based on teicoplanin therapy of staphylococcal infections; the evidence is less conclusive for vancomycin.

For both vancomycin and teicoplanin, there is data to link predose concentrations to toxicity (nephrotoxicity for vancomycin and thrombocytopenia for teicoplanin). Table 2 summarizes present recommendations for glycopeptide TDM, including those which have been used and criticized in the past, and more streamlined recommendations that may be more appropriate for the future.

Ther Drug Monit, Vol. 20, No. 5, 1998

TABLE 2. Recommendations for therapeutic drug monitoring glycopeptides

	Patient group	Serum concentrations
Vancomycin	All patients on therapy >4 to 5 days (>2 days if conservative)	Pre-dose only, range 5-15 mg/L; Predose, 5-10 mg/L; post-dose 20-40 mg/L. If conservative
Teicoplanin	Severe infection	Predose, >10 mg/L
	Severe infection	Predose, >10 mg/L (>20 mg/L if conservative)
IVDA	Severe infection	Predose, >20 mg/L
	Severe infection	Predose, >10 mg/L; post-dose, >40 mg/L
IVDA	Severe infection	Predose, >10 mg/L
	Severe infection	Predose, >10 mg/L; post-dose, >40 mg/L
IVDA	Severe infection	Predose, >10 mg/L
	Severe infection	Predose, >10 mg/L; post-dose, >40 mg/L

IVDA, Infective endocarditis; IVDA, intravenous drug abuse.

## REFERENCES

1. Gould IM. Pharmacokinetics and the relationship between in vivo and in vitro activity of antimicrobial agents. *J Chemother* 1997;9:74-83.
2. Bundran RW, Gerber AU, Holm DL, et al. Post antibiotic suppression of bacterial growth. *Rev Infect Dis* 1981;3:29-37.
3. MacKenzie PM, Gould IM. The post antibiotic effect. *J Antimicrob Chemother* 1993;32:519-37.
4. Hachinger H, Nibbeli L, Nibbeli M, et al. Post antibiotic effect of beta lactams antibiotics on Gram-negative bacteria in relation to pharmacology, initial killing and MIC. *Eur J Clin Microbiol Infect Dis* 1991;10:927-34.
5. Hachinger H. Pharmacodynamic effects of antibiotics: Studies on bacterial morphology, initial killing, post antibiotic effect and effective regrowth time. *Second J Infect Dis* 1992;1:3-52.
6. Teraoka K, Kuchi K, Shimizu K, et al. Post antibiotic effect and clinical evaluation of teicoplanin. *Nippon Kagaku Ryoku Gakki Zasshi* 1993;11:177-7.
7. Cooper MA, Yin YF, Ashby JP, et al. In vitro comparison of the post-antibiotic effect of vancomycin and teicoplanin. *J Antimicrob Chemother* 1996;26:203-7.
8. Odenbros-Torvik I, Lørdal E, Cam O. Post antibiotic sub-MIC effects of vancomycin, rifampicin, sparfloxacin and amikacin. *Antimicrob Agents Chemother* 1992;36:1832-8.
9. Latham AJ, Walker RJ, Radford JK, et al. The concentration-independent effect of amoxycillin and dicloxacillin decay in vancomycin concentrations on the killing of *Staphylococcus aureus* under aerobic and anaerobic conditions. *J Antimicrob Chemother* 1996;38:189-97.
10. Bailey EM, Rybak MJ, Kozm GW. Comparative effect of protein binding on the killing activities of teicoplanin and vancomycin. *Antimicrob Agents Chemother* 1991;35:1089-92.
11. Comanich A, Joly V, Abel L, et al. The pharmacokinetics and extracellular diffusion of teicoplanin in rabbits and comparative efficacy with vancomycin in an experimental endocarditis model. *J Antimicrob Chemother* 1982;11:321-31.
12. Chambers HF, Kennedy K. Effect of dosage, peak and trough concentrations in serum, protein binding and bactericidal rate on efficacy of teicoplanin in a rabbit model with endocarditis. *Antimicrob Agents Chemother* 1990;34:510-4.
13. Knudsen JD, Rasmussen K, Espersen F, et al. Activities of vancomycin and teicoplanin against penicillin-resistant pneumococci: in vitro and in vivo correlation to pharmacokinetic parameters in the mouse peritonitis model. *Antimicrob Agents Chemother* 1997;41:1910-3.
14. Duffet SB, Begg EJ, Chambers ST, et al. Efficacy of different vancomycin dosing regimens against *Staphylococcus aureus* determined with dynamic in vitro model. *Antimicrob Agents Chemother* 1994;38:2480-2.
15. MacGowan AP, Bowler KE. Pharmacokinetics of antimicrobial agents and rationale for their dosing. *J Chemother* 1997;9:64-73.
16. Moellering RC, Kroeger DJ, Greenblatt DJ. Pharmacokinetics of vancomycin in normal subjects and in patients with reduced renal function. *Rev Infect Dis* 1981;3:5230-3.
17. Mazza GR, McCarty RW, Halsethson CP, et al. Pharmacokinetics of vancomycin in patients with various degrees of renal function. *Antimicrob Agents Chemother* 1984;23:433-7.
18. Cunha BA, Quinlan R, Deglin BJ, et al. Pharmacokinetics of vancomycin in patients. *Rev Infect Dis* 1981;3:5269-72.
19. Blom RA, Bower LA, Miller DD, et al. Vancomycin pharmacokinetics in normal and morbidly obese subjects. *Antimicrob Agents Chemother* 1982;21:573-80.
20. Vance-Bryan K, Guss DRP, Gifford SS, et al. Effect of obesity on vancomycin pharmacokinetics: parameters as determined by using a Bayesian forecasting technique. *Antimicrob Agents Chemother* 1993;37:436-40.
21. Brown N, Ho DHW, Fong KLL, et al. Effect of hepatic function on vancomycin clinical pharmacology. *Antimicrob Agents Chemother* 1983;23:603-9.
22. Magenta RE, Arroyo JC, Ruzansky SJ, et al. Vancomycin pharmacokinetics in patients with peritonitis on peritoneal dialysis. *Antimicrob Agents Chemother* 1983;23:710-4.
23. Blarney RD, Halsethson CE, Salem NQ, et al. Pharmacokinetics of vancomycin in patients undergoing continuous ambulatory peritoneal dialysis. *Antimicrob Agents Chemother* 1984;23:603-6.
24. Touchette MA, Patel RV, Anand JV, et al. Vancomycin removal by high-flux polysulfone haemodialysis membrane in critically ill patients with end-stage renal disease. *Am J Kidney Dis* 1993;26:469-74.
25. Ureschi F, Kinowski R, de la Coma JP, et al. Clinical pharmacokinetics during continuous haemodialysis. *Clin Pharmacokinet* 1994;26:571-71.
26. Alwafel I, Najjar TA, et al. Comparison of the effects of three haemodialysis membranes on vancomycin disposition. *Am J Nephrol* 1994;24:223-8.
27. Sastre C, Levoy Q, Simon JM, et al. Pharmacokinetics of vancomycin during continuous haemodialysis. *Intensive Care Med* 1993;19:347-50.
28. Le Normand Y, Milpied N, Kerguelis MF, et al. Pharmacokinetic parameters of vancomycin for therapeutic regimen in end-stage renal patients. *Dr J Biomed Comput* 1994;36:121-3.
29. Chang D. Influence of malnutrition on the pharmacokinetics of vancomycin in infants and children. *Pediatr Infect Dis J* 1993;14:667-73.
30. Soto RB, Brundage RC, Jones PD, et al. Population pharmacokinetics of vancomycin in neonates. *Clin Pharmacol Ther* 1994;56:169-75.
31. Rotzke JC, Cronley K, Zacks D, et al. Pharmacokinetics of vancomycin: Observations in 28 patients and dosage recommendations. *Antimicrob Agents Chemother* 1983;22:191-4.
32. Pou L, Rowell M, Lopez R, et al. Changes in vancomycin pharmacokinetics during treatment. *Ther Drug Monit* 1996;18:149-53.
33. Smith JA, Thompson GA, Kenny MT, et al. Applicability of teicoplanin dosage adjustment guidelines for renally impaired patients over the range 3 to 30 mL/min. *Biopharm Drug Dispos* 1992;13:571-91.
34. Derbyshire N, Webb DB, Roberts D, et al. Pharmacokinetics of teicoplanin in subjects with varying degrees of renal function. *J Antimicrob Chemother* 1989;23:669-73.
35. Hoffler DK, Joseph P, Nussbaum E, et al. Pharmacokinetics of teicoplanin haemodialysis patients. *Infection* 1991;19:324-7.
36. Killalee-Bury D, Peyriere H, Lohjola P, et al. Influence of arteriovenous haemodialysis on teicoplanin elimination. *Br J Clin Pharmacol* 1995;40:95-7.

## PHARMACODYNAMICS, PHARMACOKINETICS, AND TDM OF GLYCOPEPTIDES

477

37. Kacet N, Dubois JP, Roussel-Devalles M, et al. Telicoplanin and amikacin in neonates with staphylococcal infection. *Pediatr Infect Dis J* 1993;12:510-3.
38. Rybak MJ, Lerner SA, Levine DP, et al. Telicoplanin pharmacokinetics in intravenous drug abusers being treated for bacterial endocarditis. *Antimicrob Agents Chemother* 1991;33:696-700.
39. Steer J, Papini R, Wilson APR, et al. Pharmacokinetics of a single dose of telicoplanin in hemiparesis. *J Antimicrob Chemother* 1996;47:545-53.
40. Jortholay O, Tob M, Rizzo N, et al. Population pharmacokinetic study of telicoplanin in severely neutropenic patients. *Antimicrob Agents Chemother* 1996;40:1243-7.
41. MacGowan AP, MacMullen CM, White LO, et al. Serum monitoring telicoplanin. *J Antimicrob Chemother* 1992;30:399-402.
42. Edwards DJ, Penco S. Routine monitoring of serum vancomycin concentrations: Waiting for proof of its value. *Clin Pharmacol* 1987;6:652-4.
43. Rodvold KA, Zafra H, Renschler JC. Routine monitoring of serum vancomycin concentrations: Can waiting be justified? *Clin Pharmacol* 1987;6:655-8.
44. Freeman CD, Quintiliani R, Nightingale CH. Vancomycin therapeutic drug monitoring: Is it necessary? *Ann Pharmacol* 1993;27:594-6.
45. Canis TO, Yamamoto-Yuue Na, Lehman PS. Serum vancomycin concentrations: Reappraisal of their clinical value. *Clin Infect Dis* 1994;18:530-43.
46. Moellering RC. Editorial: Monitoring serum vancomycin levels: Climbing the mountain because it is there. *Clin Infect Dis* 1994;18:544-5.
47. Fisher BF, Moellering RC. Retrospective study of the toxicity of preparations of vancomycin from 1974 to 1981. *Antimicrob Agents Chemother* 1983;23:138-41.
48. Welby TE, Cope AE. Impact of vancomycin therapeutic drug monitoring on patient care. *Ann Pharmacother* 1994;28:1335-9.
49. Fernandez de Gama MD, Calvo MV, Hernandez JM, et al. Cost effectiveness analysis of serum vancomycin concentration monitoring in patients with hematologic malignancy. *Clin Pharmacol Ther* 1996;60:333-40.
50. Zinnerman AE, Karon BG, Plankner KL. Association of vancomycin serum concentrations with outcome in patients with Gram-positive bacteremia. *Pharmacotherapy* 1995;15:85-91.
51. Rybak MJ, Cepulsky MJ, Rattling RC, et al. Influence of vancomycin serum concentrations on the outcome of patients being treated for Gram-positive infections. *Abstr 37th Internat Conf Antimicrob Agents Chemother* 1997:A46.
52. Wilson APR, Grunberg RN. Safety. In: *Telicoplanin: The first decade*. Abingdon, UK: The Medicine Group 1997:137-44.
53. Lepout C, Perronne C, Massip P, et al. Evaluation of telicoplanin for treatment of endocarditis caused by Gram-positive cocci. *Antimicrob Agents Chemother* 1989;33:871-6.
54. Grunberg RN. Treatment of bone, joint and vascular-associated Gram-positive bacterial infections with telicoplanin. *Antimicrob Agents Chemother* 1991;35:2392-7.
55. Rybak MJ, Cepulsky DM, Kang SL, et al. Pharmacodynamic evaluation of telicoplanin versus vancomycin in the treatment of Gram-positive bacteremia and endocarditis. *Abstr 36th Internat Conf Antimicrob Agents Chemother* 1996:A36.
56. MacGowan A, White L, Reeves D, et al. Retrospective review of serum telicoplanin concentrations in clinical trials and their relationship to clinical outcome. *J Infect Chemother* 1996;2:197-208.
57. MacGowan AP, White LO, Reeves DS, et al. Telicoplanin in *Staphylococcus aureus* septicemia: Relationship between trough serum levels and outcome. *37th Internat Conf Antimicrob Agents Chemother* 1997:A43.
58. Wilson APR, Grunberg RN, Nien H. A critical review of the dosage of telicoplanin in Europe and the USA. *Int J Antimicrob Agents* 1994;4:31-39.
59. Mulhern JL, Boocka GL, O'Shea MH, et al. Trough serum vancomycin levels predict the relapse of gram-positive peritonitis in peritoneal dialysis patients. *Am J Kidney Dis* 1993;25:611-3.

Ther Drug Monit, Vol. 20, No. 5, 1998

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.** **IMAGES ARE**  
**As rescanning these documents will not correct the image.**  
**problems checked, please do not report these problems to**  
**the IFW Image Problem Mailbox.** **the IFW Image**